ANNEX 4 IPCC Reference Approach for Estimating CO₂ Emissions from Fossil Fuel Combustion

It is possible to estimate carbon dioxide (CO₂) emissions from fossil fuel consumption using alternative methodologies and different data sources than those described in Annex 2.1. For example, the UNFCCC reporting guidelines request that countries, in addition to their "bottom-up" sectoral methodology, complete a "top-down" Reference Approach for estimating CO₂ emissions from fossil fuel combustion. Section 1.3 of the *Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories: Reporting Instructions* states, "If a detailed, Sectoral Approach for energy has been used for the estimation of CO₂ from fuel combustion you are still asked to complete...the Reference Approach...for verification purposes" (IPCC/UNEP/OECD/IEA 1997). This reference method estimates fossil fuel consumption by adjusting national aggregate fuel production data for imports, exports, and stock changes rather than relying on end-user consumption surveys. The basic principle is that once carbon-based fuels are brought into a national economy, they are either saved in some way (e.g., stored in products, kept in fuel stocks, or left unoxidized in ash) or combusted, and therefore the carbon in them is oxidized and released into the atmosphere. Accounting for actual consumption of fuels at the sectoral or sub-national level is not required. The following discussion provides the detailed calculations for estimating CO₂ emissions from fossil fuel combustion from the United States using the IPCC-recommended Reference Approach.

Step 1: Collect and Assemble Data in Proper Format

To ensure the comparability of national inventories, the IPCC has recommended that countries report energy data using the International Energy Agency (IEA) reporting convention. National energy statistics were collected in physical units from several EIA documents in order to obtain the necessary data on production, imports, exports, and stock changes.

It was necessary to make a number of modifications to these data to generate more accurate apparent consumption estimates of these fuels. The first modification adjusts for consumption of fossil fuel feedstocks accounted for in the Industrial Processes chapter, which include the following: unspecified coal for coal coke used in iron and steel production; natural gas used for ammonia production; petroleum coke used in the production of aluminum, ferroalloys, titanium dioxide, and ammonia; and other oil and residual fuel oil used in the manufacture of carbon black. The second modification adjusts for the fact that EIA energy statistics include synthetic natural gas in both coal and natural gas data. The third modification adjusts for the inclusion of ethanol in motor gasoline statistics. Ethanol is a biofuel, and it is assumed that no net CO₂ emissions occur due to its combustion. The fourth modification adjusts for consumption of bunker fuels, which refer to quantities of fuels used for international transportation estimated separately from U.S. totals. The fifth modification consists of the addition of U.S. territories data that are typically excluded from the national aggregate energy statistics. The territories include Puerto Rico, U.S. Virgin Islands, Guam, American Samoa, Wake Island, and U.S. Pacific Islands. These data, as well as the production, import, export, and stock change statistics, are presented in Table A-201.

The carbon content of fuel varies with the fuel's heat content. Therefore, for an accurate estimation of CO_2 emissions, fuel statistics were provided on an energy content basis (e.g., Btu or joules). Because detailed fuel production statistics are typically provided in physical units (as in Table A-201 for 2004), they were converted to units of energy before CO_2 emissions were calculated. Fuel statistics were converted to their energy equivalents by using conversion factors provided by EIA. These factors and their data sources are displayed in Table A-202. The resulting fuel type-specific energy data for 2004 are provided in Table A-203.

Step 2: Estimate Apparent Fuel Consumption

The next step of the IPCC Reference Approach is to estimate "apparent consumption" of fuels within the country. This requires a balance of primary fuels produced, plus imports, minus exports, and adjusting for stock

changes. In this way, carbon enters an economy through energy production and imports (and decreases in fuel stocks) and is transferred out of the country through exports (and increases in fuel stocks). Thus, apparent consumption of primary fuels (including crude oil, natural gas liquids, anthracite, bituminous, subbituminous and lignite coal, and natural gas) can be calculated as follows:

Flows of secondary fuels (e.g., gasoline, residual fuel, coke) should be added to primary apparent consumption. The production of secondary fuels, however, should be ignored in the calculations of apparent consumption since the carbon contained in these fuels is already accounted for in the supply of primary fuels from which they were derived (e.g., the estimate for apparent consumption of crude oil already contains the carbon from which gasoline would be refined). Flows of secondary fuels should therefore be calculated as follows:

Note that this calculation can result in negative numbers for apparent consumption of secondary fuels. This result is perfectly acceptable since it merely indicates a net export or stock increase in the country of that fuel when domestic production is not considered.

Next, the apparent consumption and secondary consumption need to be adjusted for feedstock uses of fuels accounted for in the Industrial Processes chapter, international bunker fuels, and U.S. territory fuel consumption. Bunker fuels and feedstocks accounted for in the Industrial Processes chapter are subtracted from these estimates, while fuel consumption in U.S. territories is added.

The IPCC Reference Approach calls for estimating apparent fuel consumption before converting to a common energy unit. However, certain primary fuels in the United States (e.g., natural gas and steam coal) have separate conversion factors for production, imports, exports, and stock changes. In these cases, it is not appropriate to multiply apparent consumption by a single conversion factor since each of its components have different heat contents. Therefore, United States fuel statistics were converted to their heat equivalents before estimating apparent consumption. Results are provided in Table A-202.

Step 3: Estimate Carbon Emissions

Once apparent consumption is estimated, the remaining calculations are similar to those for the "bottom-up" Sectoral Approach (see Annex 2.1). Potential CO_2 emissions were estimated using fuel-specific carbon coefficients (see Table A-203).¹ The carbon in products from non-energy uses of fossil fuels (e.g., plastics or asphalt) was then estimated and subtracted (see Table A-205). This step differs from the Sectoral Approach in that emissions from both fuel combustion and non-energy uses are accounted for in this approach. Finally, to obtain actual CO_2 emissions, net emissions were adjusted for any carbon that remained unoxidized as a result of incomplete combustion (e.g., carbon contained in ash or soot).²

Step 4: Convert to CO₂ Emissions

Because the IPCC reporting guidelines recommend that countries report greenhouse gas emissions on a full molecular weight basis, the final step in estimating CO_2 emissions from fossil fuel consumption was converting from units of carbon to units of CO_2 . Actual carbon emissions were multiplied by the molecular-to-atomic weight ratio of CO_2 to carbon (44/12) to obtain total carbon dioxide emitted from fossil fuel combustion in teragrams (Tg). The results are contained in Table A-204.

¹ Carbon coefficients from EIA were used wherever possible. Because EIA did not provide coefficients for coal, the IPCC-recommended emission factors were used in the top-down calculations for these fuels. See notes in Table A-204 for more specific source information.

 $^{^2}$ For the portion of carbon that is unoxidized during coal combustion, the IPCC suggests a global average value of 2 percent. However, because combustion technologies in the United States are more efficient, the United States inventory uses 1 percent in its calculations for petroleum and coal and 0.5 percent for natural gas.

Comparison Between Sectoral and Reference Approaches

These two alternative approaches can both produce reliable estimates that are comparable within a few percent. Note that the reference approach *includes* emissions from non-energy uses. Therefore, these totals should be compared to the aggregation of fuel use and emission totals from Emissions of CO₂ from Fossil Fuel Combustion (Annex 2.1) and Carbon Emitted from Non-Energy Uses of Fossil Fuels (Annex 2.3). These two sections together are henceforth referred to as the Sectoral Approach. Other than this distinction, the major difference between methodologies employed by each approach lies in the energy data used to derive carbon emissions (i.e., the actual surveyed consumption for the Sectoral Approach versus apparent consumption derived for the Reference Approach). In theory, both approaches should yield identical results. In practice, however, slight discrepancies occur. For the United States, these differences are discussed below.

Differences in Total Amount of Energy Consumed

Table A-207 summarizes the differences between the Reference and Sectoral approaches in estimating total energy consumption in the United States. Although theoretically the two methods should arrive at the same estimate for U.S. energy consumption, the Reference Approach provides an energy total that is 0.8 percent lower than the Sectoral Approach for 2004. The greatest differences lie in lower estimates for both petroleum and natural gas consumption for the Reference Approach (1.0 percent).

There are several potential sources for the discrepancies in consumption estimates:

- Product Definitions. The fuel categories in the Reference Approach are different from those used in the Sectoral Approach, particularly for petroleum. For example, the Reference Approach estimates apparent consumption for crude oil. Crude oil is not typically consumed directly, but refined into other products. As a result, the United States does not focus on estimating the energy content of the various grades of crude oil, but rather estimating the energy content of the various products resulting from crude oil refining. The United States does not believe that estimating apparent consumption for crude oil, and the resulting energy content of the crude oil, is the most reliable method for the United States to estimate its energy consumption. Other differences in product definitions include using sector-specific coal statistics in the Sectoral Approach (i.e., residential, commercial, industrial coking, industrial other, and transportation coal), while the Reference Approach characterizes coal by rank (i.e. anthracite, bituminous, etc.). Also, the liquefied petroleum gas (LPG) statistics used in the bottom-up calculations are actually a composite category composed of natural gas liquids (NGL) and LPG.
- *Heat Equivalents*. It can be difficult to obtain heat equivalents for certain fuel types, particularly for categories such as "crude oil" where the key statistics are derived from thousands of producers in the United States and abroad.
- Possible inconsistencies in U.S. Energy Data. The United States has not focused its energy data collection efforts on obtaining the type of aggregated information used in the Reference Approach. Rather, the United States believes that its emphasis on collection of detailed energy consumption data is a more accurate methodology for the United States to obtain reliable energy data. Therefore, top-down statistics used in the Reference Approach may not be as accurately collected as bottom-up statistics applied to the Sectoral Approach.
- Balancing Item. The Reference Approach uses apparent consumption estimates while the Sectoral Approach uses reported consumption estimates. While these numbers should be equal, there always seems to be a slight difference that is often accounted for in energy statistics as a "balancing item."

Differences in Estimated CO₂ Emissions

Given these differences in energy consumption data, the next step for each methodology involved estimating emissions of CO_2 . Table A-208 summarizes the differences between the two methods in estimated carbon emissions.

As mentioned above, for 2004, the Reference Approach resulted in a 0.8 percent lower estimate of energy consumption in the United States than the Sectoral Approach. The resulting emissions estimate for the Reference Approach was 0.6 percent higher. Estimates of coal and natural gas emissions using each approach yield very similar values (within 0.8 percent), though petroleum emission estimates from the Reference Approach are slightly higher (1.4 percent) than the Sectoral Approach. Potential reasons for these differences may include:

- *Product Definitions*. Coal data is aggregated differently in each methodology, as noted above. The format used for the Sectoral Approach likely results in more accurate estimates than in the Reference Approach. Also, the Reference Approach relies on a "crude oil" category for determining petroleum-related emissions. Given the many sources of crude oil in the United States, it is not an easy matter to track potential differences in carbon content between many different sources of crude, particularly since information on the carbon content of crude oil is not regularly collected.
- Carbon Coefficients. The Reference Approach relies on several default carbon coefficients by rank provided by IPCC (IPCC/UNEP/OECD/IEA 1997), while the Sectoral Approach uses annually updated category-specific coefficients by sector that are likely to be more accurate. Also, as noted above, the carbon coefficient for crude oil is more uncertain than that for specific secondary petroleum products, given the many sources and grades of crude oil consumed in the United States.

Although the two approaches produce similar results, the United States believes that the "bottom-up" Sectoral Approach provides a more accurate assessment of CO₂ emissions at the fuel level. This improvement in accuracy is largely a result of the data collection techniques used in the United States, where there has been more emphasis on obtaining the detailed products-based information used in the Sectoral Approach than obtaining the aggregated energy flow data used in the Reference Approach. The United States believes that it is valuable to understand both methods.

References

EIA (2005a) *Monthly Energy Review, September 2005* and Unpublished Supplemental Tables on Petroleum Product detail. Energy Information Administration, U.S. Department of Energy, Washington, DC, DOE/EIA-0035(2005/09).

EIA (2005b) *Monthly Energy Review*, Energy Information Administration, U.S. Department of Energy, Washington, DC. November. DOE/EIA 0035(02)-monthly.

EIA (2005c) *Annual Energy Review 2004*. Energy Information Administration, U.S. Department of Energy, Washington, DC, DOE/EIA-0384(2004). September.

EIA (1995-2005) *Petroleum Supply Annual*, Energy Information Administration, U.S. Department of Energy, Washington, DC, Volume I. DOE/EIA-0340.

IPCC/UNEP/OECD/IEA (1997) Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories, Paris: Intergovernmental Panel on Climate Change, United Nations Environment Programme, Organization for Economic Co-Operation and Development, International Energy Agency.

SAIC (2004) Analysis prepared by Science Applications International Corporation for EPA, Office of Air and Radiation, Market Policies Branch.

USGS (1998) CoalQual Database Version 2.0, U.S. Geological Survey.

Table A-201: 2004 U.S. Energy Statistics (Physical Units)

							U.S.	
Fuel Category (Units)	Fuel Type	Production	Imports	Exports	Change	Adjustment	Bunkers	Territories
Solid Fuels (Thousand Short Tons)	Anthracite Coal	1,700		a a	а			
	Bituminous Coal	546,605	i	a a	a			
	Sub-bituminous Coal	479,634		a a	а			
	Lignite	83,540		a a		3,542		
	Coke		6,873					
	Unspecified Coal		27,280	0 47,998	(18,221)	15,857		1,783
Gas Fuels (Million Cubic Feet)	Natural Gas	19,027,423	4,258,558	854,138	110,436	236,320		24,010
Liquid Fuels (Thousand Barrels)	Crude Oil	1,983,302	3,692,063	3 9,783	54,078			
	Nat Gas Liquids and LRGs	662,151	111,710	16,649	10,196			3,484
	Other Liquids	(13,459)	361,583	3 22,976	19,177			
	Motor Gasoline	167,587	181,682	2 45,498	(3,680)	85,297		41,230
	Aviation Gasoline		119					
			765	. ,				1,717
	Jet Fuel		46,50	5 14,799			150,38	12,681
	Distillate Fuel		119,118	3 40,101	(10,270)		14,89	18,284
	idual Fuel		156,024				56,25	34,869
	Naphtha for petrochemical feedstocks		48,800		(- /			
	Petroleum Coke		11,912	2 128,034	(1,939)	13,828		
	Other Oil for petrochemical feedstocks		57,80					
	Special Naphthas		2,984					
	Lubricants		2,912					281
	Waxes		1,470					
	Asphalt/Road Oil		15,60	4 2,215				
	Still Gas		(0 0	0			
	Misc. Products		49	9 1,012	640			26,660

[a] Included in Unspecified Coal

PlatacSeumees: Solid and Gas Fuels: EIA (2005a); Liquid Fuels: EIA (1995-2005).

Res

Table A-202: Conversion Factors to Energy Units (Heat Equivalents)

					Stock			U.S.
Fuel Category (Units)	Fuel Type	Production	Imports	Exports	Change	Adjustment	Bunkers	Territories
Solid Fuels (Million Btu/Short Ton)	Anthracite Coal	22.5	1					
	Bituminous Coal	23.89)					
	Sub-bituminous Coal	17.14	1			28.16		
	Lignite	12.87	1			12.87		
	Coke		24.8	30 24.80	24.80			
	Unspecified		25.0	00 25.97	20.86	27.43		25.14
Natural Gas (BTU/Cubic Foot)		1,030	1,02	23 1,009	1,030	1,025	I	1,030
Liquid Fuels (Million Btu/Barrel)	Crude Oil	5.80	5.9	98 5.80			5.8	0 5.80
	Nat Gas Liquids and LRGs	3.72	2 3.	72 3.72			3.7	2 3.72
	Other Liquids	5.83			5.83		5.8	
	Motor Gasoline	5.22	2 5.2	22 5.22	5.22	5.22	5.2	2 5.22
	Aviation Gasoline		5.0				5.0	
	Kerosene		5.0				5.6	
	Jet Fuel		5.0	67 5.67			5.6	7 5.67
	Distillate Fuel		5.8				5.8	
	Residual Oil		6.2					
	Naphtha for petrochemical feedstocks		5.2	25 5.25	5.25		5.2	5 5.25
	Petroleum Coke		6.0					
	Other Oil for petrochemical feedstocks		5.8	3 5.83			5.8	3 5.83
	Special Naphthas		5.2				5.2	
	Lubricants		6.0				6.0	
	Waxes		5.	54 5.54			5.5	
	Asphalt/Road Oil		6.0				6.6	
	Still Gas		6.0				6.0	0 6.00
	Misc. Products		5.8		5.80		5.8	0 5.80

Data Sources: Coal and lignite production: EIA (2005c); Unspecified Solid Fuels: EIA (2005b); Coke, Natural Gas and Petroleum Products: EIA (2005a).

Table A-203: 2004 Apparent Consumption of Fossil Fuels (TBtu)

Fuel Category	Fuel Type	Production	Imports	Exports	Stock Change	Adjustment	Bunkers	U.S. Territories	Apparent Consumption
Solid Fuels	Anthracite Coal	38.4			o.iai.go		24		38.4
00	Bituminous Coal	13,058.4							13,058.4
	Sub-bituminous Coal	8,220.9				11.9			8,209.0
	Lignite	1,074.8				45.6			1,029.3
	Coke		170.5	32.7	(0.7)				138.4
	Unspecified		682.0	1,246.6	(380.1)	434.9		44.8	(574.6)
Gas Fuels	Natural Gas	19,598.2	4,356.5	861.8	113.7	242.2		24.7	22,761.7
Liquid Fuels	Crude Oil	11,503.2	22,078.5	56.7	313.7				33,211.3
•	Nat Gas Liquids and LRGs	2,465.9	416.0	62.0	38.0			13.0	
	Other Liquids	(78.4)	2,106.2	133.8	111.7				1,782.3
	Motor Gasoline	874.0	947.5	237.3	(19.2)	444.8		215.0	1,373.5
	Aviation Gasoline		0.6	0.0	0.7				(0.1)
	Kerosene		4.3	7.6	(4.0)			9.7	10.5
	Jet Fuel		263.7	83.9	7.4		852.7	71.9	(608.4)
	Distillate Fuel		693.9	233.6	(59.8)		86.8	106.5	539.8
	Residual Oil		980.9	470.8	28.7		353.7	219.2	290.4
	Naphtha for petrochemical feedstocks		256.1	0.0	(1.1)				257.2
	Petroleum Coke		71.8	771.3	(11.7)	83.3			(771.1)
	Other Oil for petrochemical feedstocks		336.7	0.0	1.4	159.9			175.3
	Special Naphthas		15.7		(1.4)5	6.6			(34.9)
	Lubricants		17.7		2.9			1.7	(74.0)
	Waxes		8.1	8.5	(0.5)				0.1
	Asphalt/Road Oil		103.5	14.7	18.8				70.1
	Still Gas		0.0_{00}^{34}	2.0 0.0	0.0				0.0
	Misc. Products		0.0	5.9	3.7			154.5	145.2
Total		56,755.3	33,510.5	4,369.6	162.3	1,479.2	1,293.2	861.1	83,822.7

Note: Totals may not sum due to independent rounding.

Table A-204: 2004 Potential Carbon Dioxide Emissions

Fuel Category	Fuel Type	Apparent Consumption (QBtu)	Carbon Coefficients (Tg Carbon/QBtu)	Potential Emissions (Tg CO ₂ Eq.)
Solid Fuels	Anthracite Coal	0.038	28.26	4.0
	Bituminous Coal	13.058	25.49	1,220.5
	Sub-bituminous Coal	8.209	26.48	797.0
	Lignite	1.029	26.30	99.3
	Coke	0.138	31.00	15.7
	Unspecified	(0.620)	25.34	(53.4)
Gas Fuels	Natural Gas	22.762	14.47	1,207.7
Liquid Fuels	Crude Oil	33.211	20.33	2,475.4
·	Nat Gas Liquids and LRGs	2.795	16.99	174.1
	Other Liquids	1.782	20.33	132.8
	Motor Gasoline	1.374	19.33	97.4
	Aviation Gasoline	(0.000)	18.87	(0.0)
	Kerosene	0.010	19.72	0.8
	Jet Fuel	(0.608)	19.33	(43.1)
	Distillate Fuel	0.540	19.95	39.5
	Residual Oil	0.290	21.49	22.9
	Naphtha for petrochemical feedstocks	0.257	18.14	17.1
	Petroleum Coke	(0.771)	27.85	(78.7)
	Other Oil for petrochemical feedstocks	0.175	19.95	12.8
	Special Naphthas	(0.035)	19.86	(2.5)
	Lubricants	(0.074)	20.24	(5.5)
	Waxes	0.000	19.81	0.0
	Asphalt/Road Oil	0.070	20.62	5.3
	Still Gas	0.000	17.51	0.0
	Misc. Products	0.145	20.33	10.8
Total				6,149.8

Data Sources: Carbon content coefficients by coal rank from USGS (1998) and SAIC (2004); Unspecified Solid Fuels, Natural Gas and Liquid Fuels: EIA (2005a). Note: Totals may not sum due to independent rounding.

Table A-205: 2004 Non-Energy Carbon Stored in Products

	Consumption	Carbon	Carbon	Fraction	Carbon
	for Non-Energy	Coefficients	Content	Sequestered	Stored (Tg
Fuel Type	Use (TBtu)	(Tg Carbon/QBtu)	(Tg Carbon)		CO ₂ Eq.)
Coal	214.3	31.00	6.6	0.10	2.44
Natural Gas	380.1	14.47	5.5	0.62	12.60
Asphalt & Road Oil	1,303.9	20.62	26.9	1.00	98.58
LPG	1,437.5	16.81	24.2	0.62	55.36
Lubricants	314.8	20.24	6.4	0.09	2.16
Pentanes Plus	156.5	18.24	2.9	0.62	6.54
Petrochemical Feedstocks	a	a	a	a	57.52
Petroleum Coke	189.4	27.85	5.3	0.50	9.67
Special Naphtha	47.2	19.86	0.9	0.62	2.15
Waxes/Misc.	a	a	a	a	1.72
Misc. U.S. Territories	a	a	a	a	1.13
Petroleum					
Total					249.9

[[]a] Values for Misc. U.S. Territories Petroleum, Petrochemical Feedstocks and Waxes/Misc. are not shown because these categories are aggregates of numerous smaller components.

Note: Totals may not sum due to independent rounding.

Table A-206: 2004 Reference Approach CO₂ Emissions from Fossil Fuel Consumption (Tg CO₂ Eq. unless otherwise noted)

	Potential	Carbon	Net	Fraction	Total
Fuel Category	Emissions	Sequestered	Emissions	Oxidized	Emissions
Coal	2,083.1	2.4	2,080.7	99.0%	2,059.9
Petroleum	2,859.0	234.8	2,624.2	99.0%	2,597.9
Natural Gas	1,207.7	12.6	1,195.1	99.5%	1,189.1
Total	6,149.8	249.9	5,899.9	-	5,846.9

Note: Totals may not sum due to independent rounding.

Table A-207: Fuel Consumption in the United States by Estimating Approach (TBtu)

Approach	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
Sectoral	69,870.9	69,479.5	71,110.2	72,901.2	74,196.2	75,174.5	77,735.3	78,683.5	79,108.3	80,413.5	82,693.3	81,486.6	82,098.6	82,738.8	84,472.9
Coal	18,074.7	17,999.9	18,191.3	18,951.9	19,046.5	19,231.5	20,151.5	20,578.5	20,817.1	20,878.9	21,833.9	21,247.2	21,207.0	21,667.5	21,912.7
Natural Gas	19,375.8	19,765.4	20,438.1	20,994.7	21,455.7	22,426.5	22,837.6	22,966.8	22,559.5	22,641.7	23,554.5	22,642.7	23,370.3	22,866.9	22,987.1
Petroleum	32,420.3	31,714.3	32,480.8	32,954.6	33,694.0	33,516.5	34,746.2	35,138.2	35,731.6	36,892.9	37,305.0	37,596.7	37,521.3	38,204.4	39,573.1
Reference															
(Apparent)	69,125.2	68,348.8	69,894.8	71,695.3	73,205.3	74,113.3	76,558.8	77,991.3	78,072.3	79,240.4	81,675.2	80,741.0	81,582.5	82,149.4	83,822.7
Coal	17,602.8	17,400.9	17,725.1	18,260.8	18,724.4	18,610.6	19,519.0	20,160.7	20,033.0	20,080.7	21,039.2	20,781.3	20,875.0	21,178.1	21,898.9
Natural Gas	19,746.7	19,765.4	20,425.2	20,980.8	21,459.9	22,411.9	22,816.2	22,951.1	22,531.1	22,635.1	23,598.7	22,659.2	23,360.9	22,860.2	22,761.7
Petroleum	31,775.6	31,182.4	31,744.5	32,453.7	33,021.0	33,090.8	34,223.6	34,879.5	35,508.3	36,524.7	37,037.3	37,300.5	37,346.6	38,111.1	39,162.1
Difference	-1.1%	-1.6%	-1.7%	-1.7%	-1.3%	-1.4%	-1.5%	-0.9%	-1.3%	-1.5%	-1.2%	-0.9%	-0.6%	-0.7%	-0.8%
Coal	-2.6%	-3.3%	-2.6%	-3.6%	-1.7%	-3.2%	-3.1%	-2.0%	-3.8%	-3.8%	-3.6%	-2.2%	-1.6%	-2.3%	-0.1%
Natural Gas	1.9%	0.0%	-0.1%	-0.1%	0.0%	-0.1%	-0.1%	-0.1%	-0.1%	0.0%	0.2%	0.1%	0.0%	0.0%	-1.0%
Petroleum	-2.0%	-1.7%	-2.3%	-1.5%	-2.0%	-1.3%	-1.5%	-0.7%	-0.6%	-1.0%	-0.7%	-0.8%	-0.5%	-0.2%	-1.0%

^{*} Includes U.S. territories. Does not include international bunker fuels.

Note: Totals may not sum due to independent rounding.

Table A-208: CO₂ Emissions from Fossil Fuel Combustion by Estimating Approach (Tg CO₂ Eq.)

Approach	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
Sectoral	4,813.3	4,779.5	4,876.0	5,006.0	5,082.5	5,129.1	5,315.8	5,381.9	5,424.2	5,502.6	5,674.0	5,617.6	5,637.8	5,704.2	5,809.6
Coal	1,684.1	1,678.0	1,696.0	1,767.5	1,777.3	1,796.5	1,882.1	1,922.8	1,945.3	1,951.3	2,040.6	1,985.6	1,982.0	2,025.2	2,049.4
Natural Gas	1,013.1	1,034.0	1,070.5	1,099.0	1,121.7	1,172.6	1,194.0	1,200.3	1,177.9	1,181.6	1,228.7	1,182.0	1,219.8	1,192.9	1,198.7
Petroleum	2,116.1	2,067.6	2,109.5	2,139.4	2,183.5	2,160.1	2,239.7	2,258.7	2,301.1	2,369.7	2,404.7	2,450.0	2,436.1	2,486.1	2,561.5
Reference															
(Apparent)	4,771.6	4,717.7	4,819.9	4,951.9	5,048.9	5,092.9	5,273.5	5,383.1	5,392.4	5,455.9	5,642.1	5,608.0	5,657.9	5,719.3	5,846.9
Coal	1,641.6	1,625.0	1,655.1	1,708.1	1,752.6	1,743.4	1,827.8	1,888.8	1,880.8	1,889.0	1,977.1	1,954.6	1,963.5	1,990.8	2,059.9
Natural Gas	1,034.0	1,035.3	1,070.9	1,099.6	1,123.2	1,173.2	1,194.0	1,200.8	1,177.3	1,182.2	1,232.0	1,183.1	1,220.2	1,194.3	1,189.1
Petroleum	2,096.0	2,057.4	2,093.9	2,144.2	2,173.2	2,176.3	2,251.7	2,293.4	2,334.3	2,384.7	2,433.0	2,470.4	2,474.2	2,534.2	2,597.9
Difference	-0.9%	-1.3%	-1.2%	-1.1%	-0.7%	-0.7%	-0.8%	0.0%	-0.6%	-0.8%	-0.6%	-0.2%	0.4%	0.3%	0.6%
Coal	-2.5%	-3.2%	-2.4%	-3.4%	-1.4%	-3.0%	-2.9%	-1.8%	-3.3%	-3.2%	-3.1%	-1.6%	-0.9%	-1.7%	0.5%
Natural Gas	2.1%	0.1%	0.0%	0.1%	0.1%	0.1%	0.0%	0.0%	0.0%	0.0%	0.3%	0.1%	0.0%	0.1%	-0.8%
Petroleum	-0.9%	-0.5%	-0.7%	0.2%	-0.5%	0.7%	0.5%	1.5%	1.4%	0.6%	1.2%	0.8%	1.6%	1.9%	1.4%

⁺ Does not exceed 0.05%.

Note: Totals may not sum due to independent rounding. Includes U.S. territories. Does not include emissions from international bunker fuels.

⁺ Does not exceed 0.05%.